Contents

	st of Figures	
	st of Tables	
Pı	reface	xxviii
ln	troductiontroduction	I
1.	Impression Analysis and Its History	3
-•	1.1. What Is Impression Analysis?	
	1.2. Replication and Impression Analysis	
	1.2.1. What Are Pottery Impressions?	
	1.2.2. What Is the Replication-Based Approach?	
	1.3. The History of Research on Pottery Impressions.	
	1.3.1. The History of the Replication-Based Approach in Japanese Archaeology	
	1.3.2. Attitudes Towards Grain Impressions and Use of Replication in the United Kingdom and Scandinavia	
	1.3.3. Pottery Impressions as Archaeological Materials	
	1.4. Another Kind of Impression Analysis	
	1.4.1. A New Kind of Survey, Based on X-Ray	
	1.4.2. Tracing the Roots of Traceology: From Ushino's Replication-Based Approach to Third Excavation	10
	1.5. Conclusion	
2	Scientific Advances Brought About by the Impression Method	13
۷.	2.1. Description of Archaeological Eras in Japan	
	2.1.1. Chronology and Periodisation in Japanese Archaeology	
	2.1.2. Occupations and Cultural Characteristics of Each Period.	
	2.2. The History of Studies of Plant Usage Before the Emergence of the Pottery Impression Method	
	2.2.1. Studies of Seeds Excavated from the Soil of Archaeological Sites (Before the Year 2000)	
	2.2.2. Advancements in Scientific Investigation Methods and Applying the Pottery Impression	
	Method (2001 Onward)	
	2.3. New Historical Views Revealed by the Impression Method	
	2.3.1. Changes in Views of the History of Domesticated Plants	
	2.3.2. Increased Use of Impression Surveys and the Spread of Cultivated Plants in Japan and Korea	
	2.3.3. Development of Cereal Cultivation Throughout the Country (Verification of the Spread to Eastern Japan)	
	2.3.4. Contributions to the Reconstruction of Lifestyle History (Recognition of Household Pests)	19
3.	Innovations in Impression Methods	23
	3.1. The Limitations of the Replication-Based Approach	23
	3.2. Impression Detection and Recording Using X-Ray Equipment	24
	3.3. Results Using the New Method	25
	3.4. A New Impression Detection and Identification Method (the 'Kumamoto University Method')	27
4.	Another Step Forward (the Ultimate Impression Method)	31
	4.1. Weaknesses of the Impression Method and Methods of Overcoming Them (1)	
	4.1.1. Pottery Types Do Not Represent Time Periods	
	4.1.2. Methods of Overcoming the Issue (Extracting and Dating Carbon Embedded in Pottery)	
	4.1.3. Application of New Technology to Resolve Issues (Discoveries at the Etsuji and Kozako Sites)	32
	4.2. Weaknesses of the Impression Method and Methods of Overcoming Them (2)	33
	4.2.1. Impression Seeds Do Not Show True Quantities	33
	4.2.2. Methods of Overcoming This Issue (Adopting Various Scientific Methods of Analysis)	
	4.3. Establishing the Future of Archaeological Material Science Around Pottery: 'A Comprehensive	
	Analysis of Earthenware'	35
5.	Methods for Identifying Plants and Insects	37
-	5.1. Identifying Legume Species.	
	5.1.1. Past Methodologies for Identifying Legume Species	
	5.1.2. Legumes and Excavated Legume Pulses.	

	5.1.3. Identifying Pulses	
	5.2.1. Anatomical Characteristics of Insects	
	5.2.2. Classification of Larvae (Hornworm)	
	5.2.3. Classification of Cockroach Egg Capsules	
	5.2.4. Classifying the Pupae of Flies	
6.	New Facts About Prehistoric Japan Revealed by Impression Methods	57
	6.1. Origin of Household Cockroaches	
	6.1.1. Household Cockroaches in Modern Japan	57
	6.1.2. Archaeological Specimens in the Jōmon Period	57
	6.1.3. Identification of the Species	
	6.1.4. Establishment of Stable Lifestyle and Household Pests	58
	6.2. Species of Sitophilus (Sitophilus spp.)	
	6.2.1. Overview	
	6.2.2. Maize Weevil (Sitophilus zeamais Motschulsky)	59
	6.2.3. The Origin and History of the Dispersion of Sitophilus: Distribution According to Historical and	
	Archaeological Materials	
	6.3. Discovery of the Maize Weevil (Sitophilus zeamais) in China.	
	6.4. Discovery of Insect-Resistant Plants	
	6.5. Pottery Containing an Abundance of Seed or Animal Impressions	
	6.5.1. Introduction	
	6.5.2. The Characteristics of Seed and Arthropod Impressions in Clay: the Pottery Workspace	
	6.5.3. Pottery Containing an Abundance of Seeds	
	6.5.4. Characteristics of Seeds in Mixtures	
	6.5.5. Controversies and Issues of Concern 6.5.6. The Intention of Admixture	
	0.5.0. The intention of Admixture	🧷
7.	Identification of Key Impression Plants/Insects and Main Sources Used (Jōmon to Yayoi)	97
	7.1. Soybean: Glycine max (L.) Merr. subsp. Max; Japanese name: Daizu; Korean n.: 쿻, Chinese n.: 大豆	
	(Fig. 7.1–Fig. 7.9)	97
	7.1.1. Characteristics	97
	7.1.2. Seed/kernel morphology	97
	7.1.3. Archaeological specimens	98
	7.1.4. Usage	98
	7.2. Adzuki bean: Vigna angularis (Willd.) Ohwi et H. Ohashi var. angularis; Japanese name: Adzuki;	
	Korean n.: 팥; Chinese n. 紅豆 (Fig. 7.10–Fig. 7.16)	99
	7.2.1 Characteristics	
	7.2.2. Seed/kernel morphology	100
	7.2.3. Archaeological specimens	
	7.2.4. Usage	101
	7.3. Perilla (Beefsteak plant): Perilla spp.; Japanese name: Shiso (Egoma); Korean n.: 들깨속;	
	Chinese n. 紫蘇属 (Fig. 7.17–Fig. 7.25)	
	7.3.1. Characteristics	
	7.3.2. Seed/kernel morphology	
	7.3.3. Archaeological specimens	
	7.3.4. Usage	104
	7.4. Japanese barnyard millet: <i>Echinochloa utilis</i> ; Japanese name: Hie; Korean n.: 피; Chinese n. 稗	40.
	(Fig. 7.26–Fig. 7.30)	
	7.4.1. Characteristics	
	7.4.2. Seed/kernel morphology	
	7.4.3. Archaeological specimens	
	7.4.4. Usage	106
	7.5. Japanese lacquer tree: <i>Toxicodendron</i> spp., <i>Rhus</i> spp.; Japanese name: Urushi, Nurude, etc.;	104
	Korean n.: 옻나무; Chinese n.: 漆樹 (Fig. 7.31–Fig. 7.35)	
	7.5.1. Characteristics	
	7.5.2. Seed/kernel morphology	
	1.3.3. AMUIACOIOGICAI SPOCIIICIIS	10/

	7.5.4. Usage	
7.6.	. Hemp: Cannabis sativa L.; Japanese name: Asa; Korean n.: 삼; Chinese n.: 大麻 (Fig. 7.36–Fig. 7.38)	107
	7.6.1. Characteristics	107
	7.6.2. Seed/kernel morphology	108
	7.6.3. Archaeological specimens	108
	7.6.4. Usage	
7.7.	. Job's tear: Coix lacryma-jobi var. ma-yuen; Japanese name: Hatomugi; Korean n.: 율무;	
	Chinese n.: 台灣薏苡 (Fig. 7.39-Fig. 7.41)	109
	7.7.1. Characteristics	
	7.7.2. Seed/kernel morphology	
	7.7.3. Archaeological specimens	
	7.7.4. Usage	
7 8	Acorns: Quercus spp., Lithocarpus spp., Castanopsis spp.; Japanese name: Donguri; Korean n.:	107
7.0.	· Acons. <i>Quercus</i> spp., <i>Lunocurpus</i> spp., Custanopsis spp., Japanese name. Dongun, Korean II 도토리 (참나무속) ; Chinese n.: 椽子(栎属) (Fig. 7.42–Fig. 7.53)	110
	7.8.1. Characteristics	
	7.8.2. Seed/kernel morphology	
	7.8.3. Archaeological specimens	
7.0	7.8.4. Usage	111
7.9.	. Elymus: Elymus spp.; Japanese name: Ezomugi (Kamojigusa); Korean n.: 갯보리속;	114
	Chinese n.: 披碱草属 (Fig. 7.54–Fig. 7.56)	
	7.9.1. Characteristics	
	7.9.2. Seed/kernel morphology	
	7.9.3. Archaeological specimens	
	7.9.4. Usage	115
7.10.	. Bamboo grass: <i>Sasa</i> spp.; Japanese name: Sasa; Korean n.: 섬조릿대속; Chinese n.: 赤竹属	
	(Fig. 7.57–Fig. 7.61)	
	7.10.1. Characteristics	115
	7.10.2. Seed/kernel morphology	115
	7.10.3. Archaeological specimens	116
	7.10.4. Usage	116
7.11.	Burdock: Arctium lappa; Japanese name: Gobou; Korean n.: 우엉; Chinese n.: 牛蒡	
	(Fig. 7.62–Fig. 7.64)	116
	7.11.1. Characteristics	116
	7.11.2. Seed/kernel morphology	117
	7.11.3. Archaeological specimens	
	7.11.4. Usage	
7.12.	. Grape family: <i>Vitaceae</i> ; Japanese name: Budo; Korean n.: 포도과; Chinese n.: 葡萄科	
	(Fig. 7.65–Fig. 7.70)	117
	7.12.1. Characteristics	
	7.12.2. Seed/kernel morphology	
	7.12.3. Archaeological specimens	
	7.12.4. Usage	
7 13	. Bramble: <i>Rubus</i> spp.; Japanese name: Kiichigo; Korean n.: 산딸기속; Chinese n.: 悬钩子属	117
,.15.	(Fig. 7.71–Fig. 7.84)	119
	7.13.1. Characteristics	
	7.13.2. Seed/kernel morphology	
	7.13.2. Securities intorphology	
	7.13.4. Usage	
7 1 1	7.13.4. Osage Amur corktree: <i>Phellodendron amurense</i> ; Japanese name: Kihada; Korean n.: 황다랑어;	124
/.1 4 .	Chinese n.: 黄鳍金枪鱼 (Fig. 7.85–Fig. 7.87)	124
	7.14.1. Characteristics	
	7.14.2. Seed/kernel morphology	
	7.14.4 Users	
717	7.14.4. Usage	124
/.15.	Giant dogwood: Cornus macrophylla (Wall.) Soják; Japanese name: Kumanomizuki;	101
	Korean n.: 곰의말채나무, Chinese n.: 梾木 (Fig. 7.88–Fig. 7.89, 7.92)	
	7.15.1. Characteristics	
	7.15.2. Seed/kernel morphology	
	7.15.3. Archaeological specimens	124

	7.15.4. Usage	124
7.16	Giant dogwood: Cornus <i>controversa</i> Hemsl. ex Prain; Japanese name: Mizuki; Korean n.:	12 .
,	층층나무; Chinese n.: 灯台树 (Fig. 7.90-Fig. 7.92)	125
	7.16.1. Characteristics	
	7.16.2. Seed/kernel morphology.	
	7.16.3. Archaeological specimens	
	7.16.4. Usage	
7 17	. Japanese red elder: <i>Sambucus racemosa</i> L. subsp. <i>sieboldiana</i> (Miq.) H. Hara; Japanese name:	120
/.1/	Niwatoko; Korean n.: 덧나무; Chinese n.: 无梗接骨木 (Fig. 7.93–Fig. 7.97)	126
	7.17.1. Characteristics	
	7.17.1. Characteristics 7.17.2. Seed/kernel morphology	
	7.17.2. Securitar morphology	
	7.17.4. Usage	
7 10	Japanese prickly-ash: Zanthoxylum ailanthoides; Japanese name: Karasuzanshou;	12/
7.10	Korean n.: 머귀나무; Chinese n.: 食茱萸 (Fig. 7.98–Fig. 7.102)	127
	7.18.1. Characteristics	
	7.18.2. Seed/kernel morphology	
	7.18.2. Seed/kerner morphology	
7.10	7.18.4. Usage	129
/.19	. Beggar's lice: <i>Hylodesmum podocarpum</i> ; Japanese name: Nusubitohagi; Korean n.: 도둑놈의갈고리;	120
	Chinese n.: 山蚂蝗 (Fig. 7.103–Fig. 7.105)	
	7.19.1. Characteristics	
	7.19.2. Seed/kernel morphology	
	7.19.3. Archaeological specimens	
7.00	7.19.4. Usage	129
7.20	. Ox knee: Achyranthes bidentata var. japonica; Japanese name: Inokozuchi; Korean n.: 쇠무릎;	120
	Chinese n.: 牛膝 (Fig. 7.106–Fig. 7.108)	
	7.20.1. Characteristics	
	7.20.2. Seed/kernel morphology	
	7.20.3. Archaeological specimens	
7.01	7.20.4. Usage	131
7.21	. Agrimony: Agrimonia pilosa var. viscidia; Japanese name: Kinmizuhiki; Korean n.: (짚신나물속);	121
	Chinese n.: (龍牙草屬) (Fig. 7.109–Fig. 7.111)	
	7.21.1. Characteristics	
	7.21.2. Seed/kernel morphology	
	7.21.4 Users	
7.22	7.21.4. Usage	132
1.22		
	7.22.1. Characteristics	
	7.22.2. Seed/kernel morphology	
	7.22.3. Archaeological specimens	
7 22	7.22.4. Usage:	133
1.23	. Foxtail millet: Setaria italica; Japanese name: Awa; Korean n.: 조; Chinese n.: 粟	124
	(Fig. 7.117–Fig. 7.121)	
	7.23.1. Characteristics	
	7.23.2. Seed/kernel morphology	
	7.23.3. Archaeological specimens	
7.24	7.23.4. Usage	133
7.24	. Common millet: <i>Panicum miliaceum</i> ; Japanese name: Kibi; Korean n.: 기장; Chinese n.: 黍	126
	(Fig. 7.122–Fig. 7.126)	
	7.24.1. Characteristics	
	7.24.2. Seed/fruit morphology	
	7.24.4 Usage	
7.25	7.24.4. Usage	137
1.25	. Barley: <i>Hordeum vulgare</i> L.; Japanese name: Ohmugi; Korean n.: 보리; Chinese n.: 大麦	130
	(Fig. 7.127–Fig. 7.129)	
	7.25.1. Characteristics	
	7.25.2. Seed/kernel morphology	
	7.25.3. Archaeological specimens	138

7.25.4. Usage	139
7.26. Cockroach (egg capsule): <i>Periplaneta</i> spp.; Japanese name: Gokiburi; Korean n.: 바퀴벌레;	
Chinese n.: 蟑螂 (Fig. 7.130–Fig. 7.132)	139
7.26.1. Characteristics	139
7.26.2. Morphology	139
7.26.3. Archaeological specimens	140
7.27. Maize weevil: Sitophilus zeamais Motschulsky; Japanese name: Kokuzomushi; Korean n.:	
쌀바구미; Chinese n.: 米象 (Fig. 7.133–Fig. 7.140)	140
7.28. Turnip (Rape): Brassica spp.; Japanese name: Kabu (Aburana); Korean n.: 순무; Chinese n.:	
芜菁 (Fig. 7.141–Fig. 7.143)	
7.28.1. Characteristics	
7.28.2. Seed/kernel morphology	
7.28.3. Archaeological specimens	
7.28.4. Usage	142
7.29. Gourd: Lagenaria siceraria var. gourda; Japanese name: Hyoutan; Korean n.: $\stackrel{\text{\tiny 1}}{\lnot}$; Chinese n.:	
葫芦 (Fig. 7.144-Fig. 7.146)	
7.29.1. Characteristics	
7.29.2. Seed/kernel morphology	
7.29.3. Archaeological specimens	
7.29.4. Usage	143
7.30. Chestnut: Castanea crenata; Japanese name: Kuri; Korean n.: 밤나무; Chinese n.: 葫日本栗	
(Fig. 7.147–Fig. 7.149)	143
7.31. Japanese horse chestnut: Aesculus turbinata; Japanese name: Tochinoki; Korean n.: 칠엽수;	1.14
Chinese n.: 日本七叶树 (Fig. 7.150-Fig. 7.152)	144
7.32. Japanese walnut: <i>Juglans mandshurica</i> var. <i>sachalinensis</i> ; Japanese name: Onigurumi;	1.45
Korean n.: 가래나무; Chinese n.: 胡桃楸 (Fig. 7.153–Fig. 7.155)	145
7.33. Nepal hog plum: <i>Choerospondias axillaris</i> ; Japanese name: Chanchinmodoki; Chinese n.:	146
南酸枣 (Fig. 7.156—Fig. 7.158)	146
7.34. Wheat: Triticum compactum; Japanese name: Komugi; Korean n.: 밀; Chinese n.: 小麦	146
(Fig. 7.159–Fig. 7.161)	
7.34.2. Seed/kernel morphology	
7.34.4. Usage	
7.35. Buckwheat: <i>Fagopyrum esculentum</i> Moench; Japanese name: Soba; Korean n.: 메밀;	14/
Chinese n.: 蕎麥 (Fig. 7.162–Fig. 7.164)	1/19
7.35.1. Characteristics	
7.35.2. Seed/kernel morphology	
7.35.3. Archaeological specimens	
7.35.4. Usage	
7.36. Concluding Remarks	
7.50. Concluding Remarks	147
References	151
Multilingual Glossary	167
The Mark Country of the Country of t	4==
Location Map of the Main Sites Mentioned in the Book	177